



---

## Section 15.

# Predicting Residual Acceleration Effects on Space Experiments

**Emily Nelson**

**Computational Microgravity Laboratory  
M/S 105-1, NASA Glenn Research Center  
Cleveland, OH 44135**

**[Emily.S.Nelson@grc.nasa.gov](mailto:Emily.S.Nelson@grc.nasa.gov)**



## Predicting Residual Acceleration Effects on Space Experiments



---

### How can we predict residual acceleration effects?


---

Using an appropriate model of the acceleration, analysis tools include:

- ***theoretical*** analysis
  - order-of-magnitude analysis
  - exact solution of a simplified problem
  - asymptotic analysis
- ***numerical*** simulation
  - traditional finite difference/finite volume/finite element approach
  - stochastic approach
- ***experimental*** testing (ground-based)
  - ground-based facilities, e.g., KC-135, drop tower
  - vibrating platforms, centrifuge, clinostat (be sure to identify/quantify local acceleration field)
- ***examine previous experiments***/literature survey
- ***insight*** (and maybe a little luck)

## How does acceleration affect experiments?

- **Affects weight** (loading)
- **Modifies fluids transport** processes
  - natural convection
  - sedimentation, settling
  - mixing, separation
  - allows other phenomena to be unmasked through decreased convection
- **Changes stability thresholds**, e.g., interface between immiscible fluids, onset of convective instability, triggering of signal transduction pathways
- **Etc.**



1g



μg

Gravity is one type of acceleration; other accelerations can affect mass in gravity-like ways



## Predicting Residual Acceleration Effects on Space Experiments



### How can we model acceleration for analysis?

- Examine **actual data in the time domain** at or near the experiment:

$$g_i(t), \quad i = x, y, z$$

- **Separate out the various components** of residual acceleration from spectral analysis or from predictions:

- Analysis can be performed in the temporal or spectral domain
- Examine accelerations **individually**
  - **quasisteady** (<0.01 Hz): magnitude, orientation, frequency(?), duration(?)
  - **oscillatory**: frequency content, amplitudes, orientation, cutoffs, stationarity
  - **transient**: magnitude, duration, orientation, time delay between transients
- Examine accelerations **together**

One key feature:

**duration of microgravity time required**


Transformation to temporal domain



$$g_i(t) = g_{qs,i} + \sum_n g_{o,i} \sin(2\pi f_n t) + g_{t,i}(t)$$

## What can drive motion?


### Pressure gradients




And a whole host of other forces...

- mechanical stirring
- surface tension
- electromagnetic fields
- electrokinetic forces
- chemical reaction
- ...

### Boundaries





### Density gradients in continuous fluids




### Density gradients at interfaces


Particles, drops and bubbles



Immiscible fluids



Liquid bridges





## Predicting Residual Acceleration Effects on Space Experiments



### Effect of quasisteady $g$ , $g_{qs}$


QUASISTEADY G

- “Quasisteady” is (somewhat arbitrarily) defined as  $f \leq 0.01$  Hz
- Primary contributions to quasisteady accelerations are due to **atmospheric drag** and **gravity-gradient forces**
  - Drag is a function of attitude, vehicle geometry, local velocity, local density (and therefore, altitude, day/night, solar activity, ...)
  - Gravity-gradient forces increase with increasing distance from the center of mass
- Researchers must consider **experiment sensitivity** to:
  - **magnitude** of  $g$  (upper and lower thresholds) (expect a few  $\mu g$  on the Shuttle and on the International Space Station)
  - **orientation** of  $g$  (expect at least several degrees of variation in orientation over an orbital period)
  - in some cases, an experiment’s quasisteady regime may not coincide with this definition and temporal variations must be considered

## Effect of g on drops, particles and bubbles

QUASISTEADY G

$$\Sigma \tilde{F} = m\tilde{a}$$



$$\Sigma F = B - W \pm D + \dots$$

### Similarities:

- all are **discrete phases** surrounded by fluid
- all have **buoyant forces** acting on them (weight of displaced fluid)


### Differences:

- **different density ratios** w.r.t surrounding fluid  
(Drop:  $\rho_d > \rho_f$  Bubble:  $\rho_b < \rho_f$  Particle:  $\rho_p < \rho_f$  or  $\rho_p > \rho_f$  or  $\rho_p = \rho_f$ )
- **sign of drag force** will be a function of  $(\rho_f - \rho_m)$ ,  $l=b,d,p$   
(drag opposes direction of motion)
- **response to applied shear and pressure forces**  
(does it deform?)
- **mobility of surface** (can there be a velocity jump across the interface?)

Note: surface forces become more important with decreasing radius, acceleration, density variation

For further reading, see the excellent review by Michaelides (1997) and the book by Subramanian and Balasubramanian (2001)

## Equation of motion for discrete phase



Normal component of velocity of air bubbles in silicone oil near a wall on the Shuttle

- Ishikawa et al. (1994)

• In order to predict the discrete phase motion, one can employ:

- **creeping flow** assumptions
- **finite Reynolds number** approximations
- **semi-empirical** equations

(see, e.g., Michaelides, 1997; Subramanian and Balasubramanian, 2001)

• **Slow bubble drift** apparent in Shuttle data, almost certainly in response to quasisteady g (Ishikawa et al., 1994; Farris et al., 1998)


• Analysis of particle/bubble motion is **complicated by:**

- **wall effects** and
- **interactions** among bubbles/particles
- lack of correlation to measured **acceleration**

## Effect of quasisteady g on particles/bubbles

QUASISTEADY G

### Particle trajectories on the Shuttle



Polystyrene particles of 200, 400, 600  $\mu\text{m}$  in triglycerine sulfate on the Shuttle

- Sun et al. (1994)

**A mystery:** Why aren't the particles moving in the same direction?


#### Relevant studies on bubble/particle interaction and space data

Numerical/theoretical: Bunner and Tryggvason (1999, bubbles); Drolet and Viñals (1998, particle/wall); Ellison et al. (1995, particles/wall); Langbein (1991, bubbles)

Experimental: Farris et al. (1998, bubbles); Kawaji et al. (1999, bubble); Ishikawa et al. (1994, bubble/wall); Tryggvason et al. (2001, particles); Trolinger (2000, particles); Ellison et al. (1995, particles); Langbein (1991); Sun et al. (1994, particles)

## Effect of quasisteady $g$ on immiscible interfaces

QUASISTEADY  $g$




Even large droplets can be spherical in microgravity

In low  $g$ , the fluid that preferentially wets the walls will encapsulate the other fluid (to the best of its ability)

The shape of the interface at low  $g$  is a function of wetting properties, relative volumes of the fluids, chamber geometry, and  $g$

## Newton's 2nd law (conservation of momentum)

QUASISTEADY G



$$\sum \tilde{F} = m\tilde{a}$$

In the vertical direction, the dominant forces are:


$$\sum F = B - W \pm D + \dots = ma$$

**Forces**

**Reaction to forces**

March 5-7, 2002

PARTICLE / DROP / BUBBLE



CONTINUOUS FLUID

$$\rho = \frac{m}{V}$$

$$\frac{\partial}{\partial t}(\rho \tilde{u}) + \tilde{u} \cdot \nabla(\rho \tilde{u}) = \nabla \cdot (\mu \nabla \tilde{u}) - \nabla p + \rho \tilde{g} + \dots$$

**Reaction to forces**

**Forces**



# Predicting Residual Acceleration Effects on Space Experiments



## Governing equations for basic natural convection

For basic natural convection for Newtonian fluids with constant properties and no internal sources, we can write conservation of momentum, species and energy (using the Boussinesq approximation) as:

temporal change + convection = diffusion + source

momentum

$$\frac{\partial \tilde{u}}{\partial t} + \tilde{u} \cdot \nabla \tilde{u} = \nu \Delta \tilde{u} - \frac{1}{\rho_0} \nabla p + \beta \Delta T \tilde{g}$$

energy

$$\frac{\partial T}{\partial t} + \tilde{u} \nabla \cdot T = \alpha \Delta T$$

species


$$\frac{\partial C}{\partial t} + \tilde{u} \nabla \cdot C = D \Delta C$$

Applying scaling analysis to these equations make nondimensional numbers pop out

$$\text{Pr} = \frac{\nu}{\alpha} \quad \text{Sc} = \frac{\nu}{D}$$

Prandtl number    Schmidt number

## Example: natural convection in a molten semiconductor



**Velocity**

**Temperature**

**Species**


**Se in GaAs**

Ratio of momentum diffusion to thermal diffusion is small:  $Pr = \nu/\alpha = 0.01$

Ratio of momentum diffusion to species diffusion is large:  $Sc = \nu/D = 30$

## Effect of quasisteady g orientation on natural convection

QUASISTEADY G




Orientation of g can cause different flow modes with increased/decreased convective intensity and variation in far-field mixing

Other parameters: system geometry, boundary conditions, material properties, ...

- Arnold et al. (1991)

## Sensitivity of directional solidification to quasisteady g orientation

QUASISTEADY G



Be aware that any inhabited spacelab is likely to be ***extremely*** variable in  $\theta$  due to the rich variety of acceleration sources!

NOTE: For other experiments, this tendency towards improved mixing may actually be beneficial!

- Arnold et al. (1991)



# Predicting Residual Acceleration Effects on Space Experiments



## Effect of transient $g$ , $g_t$

TRANSIENT G

- Transient accelerations are of **short duration** by definition (<1 s to several seconds, typically)
- Causes are such things as: thruster firings, hab soars, and crew activity, e.g., hammering
- Effects can **dissipate with distance** from the source
- Researchers must consider effect of:
  - impulse **magnitude** and **duration** (or a combination of the two)
  - **orientation** of impulse
  - **time delay** between impulses


### Transient disturbances on the Shuttle

disturbance	rss magnitude ( $\mu g$ )	duration (s)
Thruster firing (OMS)*	20,000-50,000	<40
Thruster firing (PCRS)*	6000-55,000	0.001-30
Thruster firing (VCRS)*	300-700	<2
Crew activity (banging mallet)	2000	<1

\*NOT representative of Space Station thruster firings

## Effect of transient impulses

TRANSIENT G




Net acceleration=0, but system reacts in a *transient* manner with finite response time

⇒ **Net system response may be nonzero**

## Effect of transient pulse/antipulse (cont'd)

TRANSIENT G




**NOTE: Especially important for high Pr or Sc number flows**

- Monti et al. (1990)

$$(Pr = \nu / \alpha, Sc = \nu / D)$$


## Effect of PRCS thruster burns on directional solidification (MEPHISTO)

TRANSIENT G



**Results from MEPHISTO experiment**

- Favier et al. (1994)



**Results from MEPHISTO numerical model**

- Alexander et al. (1997)

Note: Seebeck voltage is proportional to the solid/liquid interface temperature




# Predicting Residual Acceleration Effects on Space Experiments



## Effect of VCRS burns on flame balls (SOFBALL)

TRANSIENT G



For a general discussion of g-jitter effects on combustion, start with Ross et al. (1998)

- Ronney et al. (1998)



# Predicting Residual Acceleration Effects on Space Experiments



## Effect of oscillatory g, $g_{osc}$

OSCILLATORY G

- **Rich frequency band** on ISS and Shuttle arising from structural oscillation, crew exercise, equipment operation
- Oscillatory g **will vary from lab to lab** on the ISS; it will depend on the disturbances that are **present** and the **experiment proximity**
- Researchers must consider **experiment sensitivity** to oscillatory g:
  - **particular frequencies**? Limitations on **bulk flows** generated from all of the frequency components?
  - **amplitude** of g (upper and lower thresholds)
  - **orientation** of g (expected to be highly variable due to variety of sources)

### Periodic disturbances on the Shuttle

disturbance	rms magnitude ( $\mu g$ )	frequency range (Hz)
Quasisteady acceleration	1-4	<0.01
Structural vibration	2-300	2.4, 3.6, 4.7, 5.2, 6.2, 7.4, 8.5
Crew exercise (ergometer)	50-1000	1-1.5, 2-3
Crew exercise (treadmill)	100-200	1-2
KU-band antenna	40-300	17.3
Life Sci refrigerator/freezer	300-400	15+




# Predicting Residual Acceleration Effects on Space Experiments




## Experiment response to oscillatory acceleration input

OSCILLATORY G



**liquid bridges**





**natural convection**

- For example, see Nelson (1991), Alexander et al. (1990), Benjapiyaporn et al. (2000)

## Body force vs. boundary vibration

OSCILLATORY G




- Ellison et al. (1995)

In a 2D numerical simulation of particles and liquid in a container with **flexible boundaries**, Ellison et al. (1995) found that transient bulk flows could be generated by Shuttle-type g-jitter. **Particles in the same plane moved in parallel.**

Studying fluid near a boundary, Volfson and Viñals (2001) found that **random vibration of boundaries** can lead to **diffusion layers** that are larger than that of pure sinusoidal vibration.

## Effect of oscillatory acceleration on bubbles

OSCILLATORY G



- Ishikawa et al. (1994)

Oscillatory response of a bubble in silicone oil to controlled sinusoidal forcing on the Shuttle

$$x(t) = \frac{6\nu A}{2\pi f} \sin(2\pi ft) - R^2 A \left[ \cos(2\pi ft) - \exp\left(-\frac{6\nu}{R^2} t\right) \right]$$

where  $A = \frac{2R^2 g_{osc}}{36\nu^2 + R^4 (2\pi f)^2}$

- On the Shuttle, 2-5 mm air bubbles were injected into silicone oil and subjected to a controlled sinusoidal oscillation

- Note **upward drift due to quasisteady acceleration**


- Theoretical and experimental prediction of bubble position are good. Correlation weakens when:
  - bubbles are near a **wall**
  - **more bubbles** are added to the fluid
  - bubble **size increases**

- Ishikawa et al. (1994)

- **Wall effects** on bubble motion, response to oscillatory forcing and to background g were also noted by Farris et al. (1998); also see Kawaji et al. (1999).

## Effect of vibration isolation on natural convection

OSCILLATORY G




- Nelson and Kassemi (1997)

### Idealized ISS environment:

- constructed from DAC-3 (Design Analysis Cycle #3)
- used a frequency range from 0.01 to 14 Hz for several hours of simulated  $\mu g$

Use this data to create  $g(t)$ :

$$g_i(t) = g_{qs,i} + \sum_n g_{o,i} \sin(2\pi f_n t)$$



### Reminder:


For a pure sinusoid,

$$\bar{g} = 0 \text{ but}$$

$$g_{\text{rms}} = \frac{\sqrt{2}}{2} g_{\text{osc}}$$

## Effect of vibration isolation on natural convection (cont'd)

OSCILLATORY G







## Predicting Residual Acceleration Effects on Space Experiments



### Initial transient in natural convection in enclosures: Startup of multifrequency sinusoidal disturbance


OSCILLATORY G




Concentration variation at solid/liquid interface as a function of time using a simplified spectrum of the Shuttle acceleration environment exhibits startup phenomenon

- Alexander et al. (1991)


## Effect of $g$ on tracks of *Euglena gracilis*



1g



$\mu g$



clinostat

("simulated  $\mu g$ ")

- Vogel et al. (1993)

Wiggles in clinostat traces are undoubtedly caused by variation in ***g orientation***

***Modulation in g magnitude*** should produce correlated ***modulation in velocity*** for microbes exhibiting gravikinesis



## Predicting Residual Acceleration Effects on Space Experiments



---

### Conclusions

---

Space experiments typically occur in a more complicated acceleration environment than that on earth.

- A known, steady acceleration environment is substituted for an acceleration environment that is ***not known a priori*** and varies significantly in terms of ***magnitude, orientation*** and ***frequency*** content
- More familiar phenomena driven by, e.g., buoyancy-driven convection, are ***dominated by less familiar forces***, e.g., surface tension, radiation heat transfer, wall effects, etc.

Nevertheless, there are things we can say with respect to the hydrodynamic effects of the microgravity environment and its effects on continuous fluids with density gradients and embedded discrete phases (bubbles, drops, particles) within fluids



# Predicting Residual Acceleration Effects on Space Experiments



## Nomenclature

### Roman characters

a	acceleration
$B=pgV$	buoyancy
C	concentration
$c_p$	heat capacity
D	drag
$D_C$	diffusivity of species
$D_m$	mass diffusivity
F	force
g	gravity
k	thermal conductivity
m	mass
p	pressure
Pr	Prandtl number= $v/\alpha$
S	source term
Sc	Schmidt number= $v/D$
u	velocity
V	volume
$W=mg$	weight

### Greek characters

$\alpha=k/\rho c_p$	thermal diffusivity
$\mu$	absolute viscosity
$\nu$	viscosity (momentum diffusivity)
$\rho$	density
$\sigma$	surface tension
$\tau$	shear stress. For Newtonian fluid, 2D, cartesian:

$$\tau = \mu \left( \frac{\partial \tilde{u}}{\partial x} + \frac{\partial \tilde{v}}{\partial y} \right)$$

### Subscripts/Superscripts

b	bubble
d	droplet
i	spatial index
l	species index
m	fluid medium
n	temporal index
osc	oscillatory
p	particle
qs	quasisteady
t	transient



## Predicting Residual Acceleration Effects on Space Experiments



---

### Bibliography

---

Alexander, J.I.D. "Low-gravity experiment sensitivity to residual acceleration: a review." **Microgravity Science and Technology** 3:52-68 (1990).

Alexander, J.I.D., J. Ouazzani, and F. Rosenberger. "Analysis of the low gravity tolerance of Bridgman-Stockbarger crystal growth: Part II. Transient and periodic accelerations." **J Crystal Growth** 97:285-302 (1991).

Alexander, J.I.D., A. Lizee, J.P. Garandet, and J.J. Favier. "Quantitative experimental characterization of g-jitter effects on directional solidification." **AIAA PAPER 97-0675** (1997).

Arnold, W., D. Jacqmin, R. Gaug and A. Chait. "Three-dimensional flow transport modes in directional solidification during space processing." **J Spacecraft and Rockets** 28:238-243 (1991).

Bunner, B. and G. Tryggvason. "Direct numerical simulations of three-dimensional bubbly flows." **Physics of Fluids** 11: 1967-1969 (1999).

Benjapiyaporn, C., V. Timchenko, E. Leonardi, G. de Vahl Davis and H.C. de Groh III. "Effects of space environment on flow and concentration during directional solidification." **International Journal of Fluid Dynamics**, vol 4, article 3 (2000). [[http://sibley.mae.cornell.edu/IJFD/2000\\_vol4/paper3/index.html](http://sibley.mae.cornell.edu/IJFD/2000_vol4/paper3/index.html)] Also **NASA TM-2000-209293** (2000).

Casademunt, J., W. Zhang, J. Viñals and R. F. Sekerka. "Narrow band noise as a model of time-dependent accelerations - Study of the stability of a fluid surface in a microgravity environment." **AIAA PAPER 93-0911**(1993).

De Groh, H.C. and E.S. Nelson. "On residual acceleration during space experiments." **ASME HTD-Vol 290**, pp 23-33 (1994).

Demel, K. "Implications of acceleration environments on scaling materials processing in space to production." In *Measurement and characterization of the acceleration environment on board the Space Station*. NASA/MSFC and Teledyne Brown (Aug 11-14, 1986).



## Predicting Residual Acceleration Effects on Space Experiments



---

### Bibliography (cont'd)

---

Drolet, F. and J. Viñals. "*Fluid flow induced by a random acceleration field.*" **Microgravity Science and Technology 11**:64-68 (1998).

Duval, W. M. B. and D. A. Jacqmin. "*Interfacial dynamics of two liquids under an oscillating gravitational field.*" **AIAA Journal 28**: 1933-1941 (1990).

Ellison, J., G. Ahmadi, L. Regel, and W. Wilcox. "*Particle motion in a liquid under g-jitter excitation.*" **Microgravity Science and Technology 8**:140-147 (1995).

Farris, S., K. S. Rezkallah, and J. D. Bugg. "*A study on the motion of a bubble subjected to simulated broadband g-jitter: Results from a recent shuttle flight.*" **Journal of the Japan Society for Microgravity Applications 15**:87-91 (1998).

Favier, J. J., J. P. Garandet, A. Rouzaud and D. Camel. "*Mass transport phenomena during solidification in microgravity; preliminary results of the first Mephisto flight experiment.*" **Journal of Crystal Growth 140**:237-243 (1994).

Favier, J. J. and D. Camel. "*Analytical and experimental study of transport processes during directional solidification and crystal growth.*" **Journal of Crystal Growth 79**:50-64 (1986).

Jenkins, J. and M. Louge. "*Microgravity Segregation of Energetic Grains.*" Science Requirements Document, NASA Glenn Research Center (1998).

Ishikawa, M., T. Nakamura, S. Yoda, H. Samejima and T. Goshozono. "*Responsive motion of bubbles to periodic g-jitter.*" **Microgravity Science and Technology 7**:164-168 (1994).

Kawaji, M., N. Ichikawa, A. Kariyasaki and A.B. Tryggvason. "*Large bubble motion in a fluid cell under microgravity: ISCAP experiments on the effects of g-jitter and forced vibration.*" **IAF Paper 99-J308** (1999).



## Predicting Residual Acceleration Effects on Space Experiments



---

### Bibliography (cont'd)

---

Langbein, D. "*Motion of ensembles of spherical particles in a fluid due to g-jitter.*" **Advances in Space Research** **11**:189-196 (1991).

Lizee, A. and J. I. D. Alexander. "*Chaotic thermovibrational flow in a laterally heated cavity.*" **Physical Review E** **56**: 4152-4156 (1997).

Michaelides, E.E. "*Review - The transient equation of motion for particles, bubbles, and droplets.*" **Journal of Fluids Engineering** **119**:233-247 (1997).

Meseguer, J., L.A. Slobozhanin and J. M. Perales. "*A review on the stability of liquid bridges.*" **Advances in Space Research** **16**:5-7 (1995).

Monti, R. "*Gravity jitters: effects on typical fluid science experiments.*" In J.N. Koster and R.L. Sani, *Low-gravity fluid dynamics and transport phenomena.* AIAA (1990).

Nelson, E.S. "*An examination of anticipated g-jitter on Space Station and its effects on materials processes.*" **NASA TM 103775** (1991, 1994).

Nelson, E.S. and M. Kassemi. "*The effects of residual acceleration on concentration fields in directional solidification.*" **AIAA 97-1002** (1997).

Ronney, P.D., M-S Wu, H.G. Pearlman, and K.J. Weiland. "*Structure of Flame Balls At Low Lewis-number (SOFBALL) - Preliminary results from the STS-83 and STS-94 space flight experiments.*" **AIAA Paper 98-0463** (1998).

Ross, H. D. "*A short collation of g-jitter effects on combustion systems with realistic g-jitter.*" Paper Number 21, 17th International Microgravity Measurements Group Meeting. Cleveland, OH (1998).



## Predicting Residual Acceleration Effects on Space Experiments



---

### Bibliography (cont'd)

---

Subramanian, R.S. and R. Balasubramanian. "*The motion of bubbles and drops in reduced gravity.*" Cambridge University Press (2001).

Sun, J., F. M. Carlson, L. L. Regel, W. R. Wilcox, R. B. Lal and J. D. Trolinger. "*Particle motion in the fluid experiment system in microgravity.*" **Acta Astronautica** 34:261-269 (1994).

Trolinger, J. D., R. H. Rangel, C. F. M. Coimbra, R. B. Lal, W. Witherow and J. Rogers. "*SHIVA: Spaceflight Holography Investigation in a Virtual Apparatus.*" (2000).

Tryggvason, B.V., R.F. Redden, R.A. Herring, W.M.B. Duval, R.W. Smith, K.S. Rezkallah, and S. Varma. "*The vibration environment on the International Space Station: Its significance to fluid-based experiments.*" **Acta Astronautica** 48:59-70 (2001).

Vogel, K., R. Hemmersbach-Krause, C. Kühnel and D.-P. Häder. "*Swimming behavior of the unicellular flagellate, Euglena gracilis, in simulated and real microgravity.*" **Microgravity Science and Technology** 5:232-237 (1993).

Volfson, D. and J. Viñals. "*Flow induced by a randomly vibrating boundary.*" **Journal of Fluid Mechanics** 432:387-408 (2001).